CLINICAL GUIDELINES ID TAG		
Title:	Guideline for anaesthetic management of fractured neck of femur	
Author:	Dr Nidhi Gupta	
Speciality / Division:	Anaesthetics / ATICS	
Directorate:	Acute	
Date Uploaded:	11 th January 2017	
Review Date	10 th January 2020	
Clinical Guideline ID	CG0397	

Guidelines for Anaesthetic Management of fracture NOF (Cannulation screws, Hemi, DHS, Gamma Nail) For THR-follow Enhanced RecoveryTHR guidelines

PRE- OP	San A. 7 of transma for management of outin	as and ation and as manhidities	
PRE- OP	See A-Z of trauma for management of anticoagulation and co-morbidities Restrict fasting to Solids (2 am for AM list and 7am for PM list) Liquids (6 am for AM list and 11am for PM list) Continue transdermal analgesic patches Pre-op drinks as per agreed protocol		
	Application of Bair hugger 2 hours prior to theatre		
INTRA-OP	Spinal preferential to GA unless contraindicated		
	Administer nerve block (Fascia iliaca, femoral nerve or lumbar plexus) before positioning for spinal or after spinal or GA If no nerve block performed then administer LIA as per THR protocol		
	Low threshold for invasive arterial monitoring esp in high risk patients		
	Low dose spinal <10mg 0.5% plain	GA and IV morphine	
	bupivacaine	C-1 mile 2 × miles parameter	
	Dilute with 0.9% NaCl to increase volum	e	
	to 5ml if necessary		
	No IT diamorphine		
	If IT opioid required <25 mcg fentanyl		
	Avoid sedation where possible. Propofol agent of choice if required		
CI: PE, previous DVT, CVA, caution in stents			
	IV magnesium infusion 3g in females and 4g in males		
	Maintain systolic BP within 25% of pre-induction or pre-spinal level Maintain MAP > 65mmHg		
CEMENTING	Be aware of Bone cement implantation syndrome Discuss possibility of non-pressurised cementing if high risk patient with the surgeon		
POST-OP	Day 0 – 2	Day 0 – 2	
Regular	Paracetamol 1g IV 6hrly for 24 hrs then PO	Paracetamol 1g IV 6hrly for 24 hrs then PO	
	Codeine 30-60 mg PO 6 hrly (if needed)	Codeine 30-60 mg PO 6hrly (if needed)	
Rescue	Shortec 5-10mg PO 2-4 hrly PRN	Shortec 5-10mg PO 2-4 hrly PRN	
	Day 3	Day 3	
Regular	Paracetamol 1g PO 6hrly	Paracetamol 1g PO 6 hrly	
Rescue	Shortec 5-10mg PO 2-4 hrly PRN	Shortec 5-10mg PO 2-4 hrly PRN	
	Prior to leaving theatre		
	 Age and renal function adjusted post-op analgesia prescribed Post-operative fluids prescribed Antiemetics or laxatives prescribed 		